



Black hole spin measurements with NuSTAR

Andrea Marinucci
(Università degli Studi Roma Tre)
on behalf of the
NuSTAR AGN Physics WG

The 7th FER0 Meeting
Finding Extreme Relativistic Objects
Krakow, 28th-30th August 2014

Outline

- Brief introduction about scientific goals
 - Radio-quiet AGN seen by NuSTAR
 - Results
 - Conclusions



Outline

- Brief introduction about scientific goals
 - Radio-quiet AGN seen by NuSTAR
 - Results
 - Conclusions

Introduction – Primary emission



- One of the main open problem for AGN is the nature of the primary X-ray emission.

- It is due to Comptonization of soft photons, but the geometry, optical depth and temperature of the emitting corona are largely unknown.




ular models in $E_{\text{cut}} = 2-3 kT$, so measuring E_{cut} helps constrain


Introduction – Relativistic reflection



Light bending model:
much of the flux is bent onto the disk
giving a constant, strong RDC



Introduction – Relativistic reflection



Spin alters shape of Fe $K\alpha$ line and Compton hump in predictable, measurable ways.

Outline

- Brief introduction about scientific goals
 - Radio-quiet AGN seen by NuSTAR
 - Results
 - Conclusions

The NuSTAR satellite

Nuclear Spectroscopic Telescope Array

1 Ms Sensitivity

3.2×10^{-15} erg/cm²/s (6 – 10 keV)

1.4×10^{-14} erg/cm²/s (10 – 30 keV)

Imaging

HPD 58"

FWHM 18"

Localization 2" (1-sigma)

Harrison et al., 2013

Spectral response

energy range: 3-79 keV

ΔE @ 6 keV 0.4 keV FWHM


ΔE @ 60 keV 1.0 keV FWHM

Target of Opportunity

response <24 hr

typical 6-8 hours


80% sky accessibility



The NuSTAR satellite

Benefit of NuSTAR high effective area and low background yields $\sim 100\times$ better S/N versus

Swift-MCG-6-30-15: 125 ks net exposure time and same 15-70 keV flux (6.5×10^{-11} erg/cm²)



Marinucci et al., 2014a

Radio-quiet AGN observed by NuSTAR

Target	Exposure Time	Simultaneous	Reference
Ark 120	130 ks	XMM-Newton	Matt et al., 2014
IC 4329A	180 ks	Suzaku	Brenneman et al., 2014a,b
MCG—6-30-15	3x130ks	XMM-Newton	Marinucci et al., 2014a
Mrk 335	300 ks	Swift	Parker et al., 2014
NGC 1365	4x130 ks	XMM-Newton	Risaliti et al., 2013 Walton et al., 2014
SWIFT J2127.4	3x130ks	XMM-Newton	Marinucci et al., 2014b

Radio-quiet AGN observed by NuSTAR

Target	Exposure Time	Simultaneous	Reference
Ark 120	130 ks	XMM-Newton	Matt et al., 2014
IC 4329A	180 ks	Suzaku	Brenneman et al., 2014a,b
MCG—6-30-15	3x130ks	XMM-Newton	Marinucci et al., 2014a
Mrk 335	300 ks	Swift	Parker et al., 2014
NGC 1365	4x130 ks	XMM-Newton	Risaliti et al., 2013 Walton et al., 2014
SWIFT J2127.4	3x130ks	XMM-Newton	Marinucci et al., 2014b

Outline

- Brief introduction about scientific goals
 - Radio-quiet AGN seen by NuSTAR
 - **Results**
 - Conclusions


The soft excess in Ark 120

Most AGN show soft X-ray emission in excess of the extrapolation of the hard primary emission


In many sources the soft excess is well explained by ionized reflection from the accretion disk (e.g. Walton et al. 2013)

However, there are sources in which another component is required (Patrick et al. 2012, Lohfink et al. 2012, Petrucci et al. 2013)

Ark 120 is one of them (Matt et al. 2014)




Ross & Fabian 2005




Matt et al. 2014


The soft excess in Ark 120



Matt et al. 2014



No obvious evidence for a relativistic Iron line likely from a previous Suzaku observation, Nardini et



The broad-band best fit is with a Comptonization model for the soft excess. Optxagnb (Done et al. 2012) is a disk/corona emission model which assumes a thermal disk emission outside the coronal radius, and soft and hard Comptonization inside.


The soft excess in Ark 120

Matt et al. 2014

a	0	0.50	0.99
L/L_{Edd}	$0.16^{+0.16}_{-0.08}$	$0.05^{+0.01}_{-0.01}$	$0.04^{+0.03}_{-0.01}$
$R_c (R_G)$	$11.5^{+0.1}_{-3.4}$	$31.3^{+39.2}_{-16.6}$	$24.9^{+16.0}_{-15.2}$
kT (keV)	$0.33^{+0.02}_{-0.02}$	$0.32^{+0.01}_{-0.01}$	$0.32^{+0.02}_{-0.01}$
τ	$12.9^{+1.1}_{-0.9}$	$13.6^{+0.6}_{-0.2}$	$13.6^{+0.4}_{-0.7}$
Γ	$1.73^{+0.02}_{-0.02}$	$1.73^{+0.02}_{-0.02}$	$1.73^{+0.02}_{-0.02}$
E_c (keV)	>190	>190	>190



on board on XMM-Newton support an interme

n.




Relativistic reflection in MCG—6-30-15


First broad Fe Ka line ever observed (Tanaka+95) and interpreted as originating from a rapidly spinning BH (Iwasawa+96)



Relativistic reflection in MCG—6-30-15




Brenneman et al., in prep.




Miniutti et al., 2007

- Residuals to a power-law are qualitatively similar to those seen in previous epochs, as is overall flux state.

Relativistic reflection in MCG—6-30-15




Marinucci et al. 2014a




The source has been observed in a very bright and variable state in 2013 during the XMM+NuSTAR campaign (Marinucci et al. 2014a)

Relativistic reflection in MCG—6-30-15




Marinucci et al. 2014a




The different spectral shape in the time intervals considered is explained in terms of the interaction between the primary continuum and the accretion disk.

Relativistic reflection in MCG—6-30-15



Marinucci et al. 2014a

Relativistic reflection in MCG—6-30-15




Brenneman et al, in prep.


- Spin, disk inclination, iron abundance linked between different intervals.

Relativistic reflection in Mrk 335

Mrk 335 was observed by NuSTAR and Swift in a very faint state, allowing us to study the reflection properties of the source.





Parker et al. 2014



<http://www.nustar.caltech.edu/news/nustar140812>


Relativistic reflection in Mrk 335



Parker et al. 2014

When flux-resolved states are considered a maximally rotating black hole spin is measured.

Black hole spin in NGC 1365




Risaliti et al. 2013, Nature

NGC 1365: a source in which both absorption and relativistic reflection play a major role in the X-rays

The first NuSTAR published paper is the spin measurement in NGC 1365



Black hole spin in NGC 1365



Walton et al. 2014

NGC 1365 was observed by XMM and NuSTAR four times. Despite large errors, no variations in the reflected components are found, demonstrating the role of

Black hole spin in NGC 1365




Walton et al. 2014

Relativistic reflection in SWIFT J2127.4

▫ NLS1 with a relativistically broadened Fe K α emission line ($a=0.6\pm0.2$), a steep continuum ($\Gamma=2-2.4$), $E_c=30-90$ keV, $L_{bol}/L_{Edd}\sim0.18$ (Miniutti+09, Malizia+08, Panessa+11, Sanfrutos+13)


▫ It was observed simultaneously with XMM-Newton for ~ 300 ks and both a strong Compton Hump and a broad Fe K α line are present



▫ Marinucci et al. 2014b


Relativistic reflection in SWIFT J2127.4

- When a model composed of a primary continuum, relativistic and distant reflection components is applied to the data the only residuals are above ~ 25 keV



Relativistic reflection in SWIFT J2127.4


When a model composed of a primary continuum, relativistic and distant reflection components is applied to the data the only residuals are above ~ 25 keV




ser +14) allows us to measure a cutoff energy $E_c = 108 \pm 10$ keV and to infer the cont

Relativistic reflection in SWIFT J2127.4

Confidence contours: Chi-Squared



Confidence contours: Chi-Squared



□ Using compTT (Titarchuk+94) with two different geometries we get:

□ SLAB

□ $kT_e = 68^{+37}_{-32}$ keV


□ $\tau = 0.35^{+0.35}_{-0.19}$

□ SPHERE

□ $kT_e = 53^{+28}_{-26}$ keV

□ $\tau = 1.35^{+1.03}_{-0.67}$

Relativistic reflection in SWIFT J2127.4



$$a = 0.58^{+0.11}_{-0.17}$$

Marinucci et al. 2014b

Outline

- Brief introduction about scientific goals
 - Radio-quiet AGN seen by NuSTAR
 - Results
- **Conclusions**

Conclusions

- The recent NuSTAR observational campaign of Radio-quiet AGN allowed us to study:
 - PRIMARY EMISSION
 - First measurements of the coronal parameters T and τ in IC 4329A
 - Warm Comptonization in Ark 120
 - (in addition to reflection?)
 - Bringing the two pieces of information together we have an unprecedented power to constrain the geometry and environment (between a relativistic and a non-relativistic disk) of the nucleus
- REFLECTED EMISSION
- EMISSION
- EMISSION
- black hole spin measurements in a number of sources: Ark 120, MCG—6-30-15, NGC